arXiv:math/0611533v1 [math.AG] 17 Nov 2006

# Addendum to "$K3$-surfaces of genus 8 and varieties of sums of powers of cubic fourfolds"

Atanas Iliev and Kristian Ranestad

## Abstract

In this note, which is an addendum to the e-print math.AG/9810121, we prove that the variety VSP(F,10) of presentations of a general cubic form F in 6 variables as a sum of 10 cubes is a smooth symplectic 4-fold, which is deformation equivalent to the Hilbert square of a K3 surface of genus 8 but different from the family of lines on a cubic 4-fold. This provides a new geometric construction of a compact complex symplectic fourfold, different from a Hilbert square of a K3 surface, a generalized Kummer 4-fold, the variety of lines on a cubic 4-fold and the recent examples of O'Grady (see Duke Math. J. 134, no. 1 (2006), 99-137).

Eyal Markman noticed that the proof of Theorem 3.17 in the paper [2] is incomplete. The purpose of this note is to correct both the statement and the proof of that theorem.

In the paper we show that $Hilb_2S$ for a general $K3$ surface $S$ of genus 8, coincides with the variety of sums of powers $VSP(F,10)$ of a cubic fourfold $F$. On the other hand Beauville and Donagi showed that $Hilb_2S$ coincides with the variety of lines $Hilb_l(F')$ in a Pfaffian cubic fourfold $F'$. The natural polarization on $VSP(F,10)$ and on $Hilb_l(F')$ provides two polarizations $L_{vsp}$ and $L_{lines}$ on $Hilb_2S$. Now the pairs $(Hilb_2S, L_{vsp})$ and $(Hilb_2S, L_{line})$ both deform smoothly with the cubic $F$ (resp. $F'$), giving components in the moduli space of polarized hyperkahler manifolds. The general element of each component has Picard group generated by the given polarization. In the proof of Theorem 3.17 in [2] we overlooked the importance of the polarization when we concluded that the two components of the moduli space coincide. In fact we prove

**Theorem.** The two polarizations $L_{vsp}$ and $L_{lines}$ are linearly independent on $Hilb_2S$.

By Huybrechts ([1] 1.8) and Lemma 3.18 of [2] we get the

**Corollary.** The moduli spaces of $VSP(F,10)$ and of $Hilb_l(F)$ for general cubic fourfolds form two distinct components of the moduli space of polarized hyperkahler manifolds. They intersect each other transversally along a divisor representing manifolds $Hilb_2S$, for a $K3$-surface $S$ of genus 8.

**Proof**. To show that the polarizations are independent we show that the degrees of a particular rational curve with respect to these two polarizations are relatively prime. The polarization $L_{lines}$ is primitive, so this suffices to conclude.

The test curve $C_p$ is a ruling in the diagonal on $Hilb_2(S)$, i.e. the pencil of length two subschemes supported at a point $p \in S$. This pencil is of course defined by the pencil of lines through $p$ in the tangent plane to $S$ at $p$. We show that this curve has degree 5 with respect to $L_{lines}$ and degree 3 with respect to $L_{vsp}$.

Recall the setting in [2]. We fix a vector space $V \cong \mathbf{C}^6$, and consider the Grassmannians $\mathbf{G}(2,V) \subset \mathbf{P}(\wedge^2 V)$ and $\mathbf{G}(4,V)$ in the natural dual space $\mathbf{P}(\wedge^4 V)$. The surface $S$ is a linear section $S = \mathbf{P}^8 \cap \mathbf{G}(2,V)$. We denote the $\mathbf{P}^8$ that span $S$ by $\mathbf{P}^8(S)$. The cubic fourfold $F'$ is the linear section defined by $\mathbf{P}^5(S) = \mathbf{P}^8(S)^\perp$ of the Pfaffian cubic in $\mathbf{P}(\wedge^4 V)$ (the secant variety of $\mathbf{G}(4,V)$). Since $\mathbf{G}(4,V) \subset \mathbf{P}(\wedge^4 V)$ is arithmetically Gorenstein and has codimension 6, the quotient of its homogeneous coordinate ring defined by $\mathbf{P}^5(S)$ is an Artinian Gorenstein ring. The socle degree of this ring is 3. The dual socle generator is therefore a cubic form. This form defines the socalled apolar cubic fourfold $F$ associated to $S$.

There are three polarizations on $Hilb_2 S$ that show up in this setting. First, $Hilb_2 S$ is parameterized by the secant (and tangent) lines to $S$ (we may assume that $S$ contains no lines), so there is a polarization $L_{sec}$ defined by the Plücker embedding of the variety of secant lines to $S$. Secondly, there is a polarization $L_{lines}$ defined by the Plücker embedding of the variety of lines in $F'$. The third polarization $L_{vsp}$ defines the vsp-map

$$f : Hilb_2\ S \to VSP(F, 10) \subset \mathbf{G}(4, \wedge^2\ V),$$

where $F$ is the apolar cubic 4-fold to $S$. By [2] Lemmas 2.8 and 3.8, it is defined as follows: We first identify $l \in Hilb_2\ S$ with a secant or tangent line to $S$. If $l$ is a secant line, then let $\{p, q\} = l \cap S$ and let $L_p$ and $L_q$ be the corresponding lines in $\mathbf{P}(V)$. Since $S$ does not contain lines (in particular $S$ does not contain the line $< p, q >$ spanned by $p$ and $q$), the span $< L_p \cup L_q >= \mathbf{P}(U_l)$ for some 4-space $U_l \subset V$. The set $Q_l$ of all lines $L \subset \mathbf{P}(V)$ that intersect $L_p$ and $L_q$ is a smooth quadric surface $Q_l \subset \mathbf{G}(2, U_l)$, and the space $\mathbf{P}(l) = \mathbf{P}^3_l = Span\ Q_l \subset \mathbf{P}(\wedge^2\ U_l)$ is just the image $f(l)$ of $l$. If $l$ is a tangent line, the same argument goes through for a quadric surface cone $Q_l$ with vertex at $p$ spanning a 3-space $\mathbf{P}(l)$. In both cases, note that $\mathbf{P}(l)$ is the polar in $\mathbf{P}(\wedge^2\ U_l)$ of $l$ with respect to the quadric $\mathbf{G}(2, U_l)$.

**Lemma**. $C_p \cdot L_{vsp} = 3$.

**Proof**. In order to describe the restriction of the vsp-map $f$ to the curve $C_p \subset Hilb_2\ S$, note first that the map $\beta : Hilb_2\ S \to \mathbf{G}(4,V)$, $\beta : l \to U_l$ defined above, is the restriction on $Hilb_2\ S \subset \mathbf{P}(\wedge^2\ V)$ of the Cremona transformation $\beta : \mathbf{P}(\wedge^2\ V) \to \mathbf{P}(\wedge^4\ V)$ defined by the quadratic Pfaffians defining $\mathbf{G}(2,V)$, see [2]. The restriction $\beta_p$ of $\beta$ to the tangent plane $\mathbf{P}^2_p$ to $S$ at $p$ is defined by the system of all conics in $\mathbf{P}^2_p$ singular at $p$. Therefore $\beta_p$ blows up $p$ and maps the exceptional divisor $E_p$ over $p$ to a conic $q_p \subset \mathbf{G}(4,V)$. In the embedding $Hilb_2\ S \subset \mathbf{P}(\wedge^2(\wedge^2\ V))$ defined above, $C_p$ is identified with the pencil of lines through $p$ in the tangent plane $\mathbf{P}^2_p$, i.e. with the exceptional curve $E_p$. Therefore the Cremona transformation $\beta$ sends the line $C_p \subset (\mathbf{P}^2_p)^*$ to the conic $q_p \subset \mathbf{G}(4, V)$ as above.

The pencil of $\mathbf{P}(\wedge^2 U_l)$'s in $\mathbf{P}(\wedge^2 V)$ corresponding to $l \in C_p$ form a rational scroll $W$ of degree 6 parameterized by the elements $\beta(l) \in q_p$ of the conic $q_p \subset \mathbf{G}(4, V)$ (the rational scroll of $\mathbf{P}(\wedge^2 U_l)$'s corresponding to a line in $\mathbf{G}(4, V)$ has degree 3). The scroll $W$ has the point $p$ as its vertex, so $W$ contains the tangent plane $\mathbf{P}^2_p$ to $S$ at $p$, and for any $l$ the ruling $\mathbf{P}(\wedge^2 U_l) \subset W$ intersects $\mathbf{P}^2_p$ at the line $l$. Finally, for each $l \in C_p$ the space $\mathbf{P}(l) = \mathbf{P}^3_l = f(l)$ coincides with the polar 3-space to the line $l \subset \mathbf{P}(\wedge^2 U_l)$ with respect to the quadric $\mathbf{G}(2, U_l)$, i.e. the isomorphic image $f(C_p) \subset VSP(F(S), 10) \subset \mathbf{G}(4, \wedge^2 V)$ is the polar scroll to the plane $\mathbf{P}^2_p$ with respect to the quadric bundle $W \cap \mathbf{G}(2, V)$. We use the following lemma to show that the degree of this polar scroll is 3.

**Lemma** Let $X = \mathbf{P}^n_{\mathbf{P}^1}$ be a $\mathbf{P}^n$-bundle with $H$ the divisor associated to $\mathcal{O}_X(1)$ and $F$ the class of a fiber. Let $Q \subset X$ be a quadric bundle such that $Q \equiv 2H + aF$, and let $s : \mathbf{P}^1 \to X$ be a section of the projective bundle such that $s^*(\mathcal{O}_X(1))$ has degree $d$. Let $Y_s \subset X$ be the variety in $X$ of polar $\mathbf{P}^{n-1}$-spaces to $s(t)$ with respect to $Q_t$ where $t \in \mathbf{P}^1$. Then

$$Y_s \equiv H + (a + d)F.$$

**Proof**. This is straightforward computation. Let $< x_i >$ be a basis for the sections of $\mathcal{O}_X(1)$, and let $u_0, u_1$ generate $H^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(1))$. Then the equation for $Q$ is quadratic in the $x_i$ and of degree $a$ in the $u_i$. The point $s(t)$ has coordinates of degree $d$ in the $u_i$. By differentiation the polar $Y_s$ has a linear equation in the $x_i$ of degree $d + a$ in the $u_i$. Q.E.D.

In our situation we first note that the quadric bundle $Q \subset W \subset G$ has degree 10. In fact the quadric bundle of quadrics $\mathbf{G}(2, U_l)$'s in the cubic scroll of the $\mathbf{P}(\wedge^2 U_l)$'s corresponding to a line in $\mathbf{G}(4, V)$ is isomorphic to a tangent hyperplane section of a $\mathbf{G}(2, 5)$, and therefore has degree 5. Since $Q$ is the corresponding quadric bundle in the sextic scroll $W$ corresponding to a conic in $\mathbf{G}(4, V)$, it has degree 10. If $H$ is the pullback (to a desingularization of $W$) of the Plücker divisor on $\mathbf{G}(2, V)$, and if $F$ is a member of the ruling, then $Q \equiv 2H - 2F$ (on the desingularization of $W$). Let $s_0 : \mathbf{P}^1 \to W$ be the constant map to the vertex $p$. Then, by the lemma,

$$Y_{s_0} \equiv H - 2F.$$

Let $Q_0 = Q \cap Y_{s_0} \subset Y_{s_0} = X_0$ and let $s_1 : \mathbf{P}^1 \to X_0$ be the section corresponding to a line in the tangent plane that does not pass through the tangency point $p$. Applying the lemma once more we get:

$$Y_{s_1} \equiv H - F$$

on $X_0$. Thus $Y_{s_1}$ has degree

$$H^4 \cdot (H - F) \cdot (H - 2F) = 3.$$

Q.E.D.

**Lemma**. $C_p \cdot L_{lines} = 5$.

**Proof**. Let $V_p$ be the 2-space corresponding to $p \in \mathbf{G}(2, V)$. Each tangent line $l$ to $\mathbf{G}(2, V)$ at $p$ is tangent to a unique quadric $\mathbf{G}(2, U_l)$ for a 4-space $U_l$ that contains $V_p$. The hyperplanes that contain $\mathbf{G}(2, U_l)$ form a $\mathbf{P}^8(U_l)$ in the dual space $\mathbf{P}(\wedge^4 V)$, and this $\mathbf{P}^8(U_l)$ is the tangent space to $\mathbf{G}(4, V)$ at the point corresponding to $U_l$. The tangent space $\mathbf{P}^8(U_l)$ is contained in the Pfaffian cubic of secant lines to $\mathbf{G}(4, V)$, and it intersects $\mathbf{P}^5(S)$ in a line $l'$, since $\mathbf{P}^5(S)^\perp = \mathbf{P}^8(S)$ intersects $\mathbf{P}(\wedge^2 U_l) = \mathbf{P}^8(U_l)^\perp$ in a line. We want to compute the degree of the curve of such lines $l'$ in $\mathbf{P}(\wedge^4 V)$ corresponding to the pencil of tangent lines to $S$ at $p$.

Above we computed the degree of $W \subset \mathbf{P}(\wedge^2 V)$, the pencil of $\mathbf{P}(\wedge^2 U_l)$'s that intersect $\mathbf{P}^8(S)$ in a tangent line to $S$. The corresponding pencil of $\mathbf{P}^8(U_l)$'s orthogonal to the $\mathbf{P}(\wedge^2 U_l)$'s is a rational 9-fold scroll $W'$ of degree 6 in $\mathbf{P}(\wedge^4 V)$. Since $W$ is a cone, the scroll $W'$ is contained in a hyperplane. Furthermore $\mathbf{P}^5(S)$ is contained in this hyperplane. In fact, since $W \cap \mathbf{P}^8(S)$ is precisely a plane (the tangent plane to $S$) through the vertex $p$ that intersect each ruling of $W$ in a line, the intersection of each $\mathbf{P}^8(U_l)$ with $\mathbf{P}^5(S)$ is a line. The degree of $W' \cap \mathbf{P}^5(S)$ coincides with the degree of the projection of $W$ from the tangent plane. But this degree is 5, so the lemma follows. Q.E.D.

**Acknowledgement.** We thank Brendan Hassett for helpful discussions on the Fano variety of lines in a cubic fourfold.

# References

[1] Huybrechts, Daniel *Compact hyper-Kähler manifolds: basic results.* Invent. Math. **135** (1999), no. 1, 63–113.

[2] Iliev, Atanas; Ranestad, Kristian $K3$ *surfaces of genus 8 and varieties of sums of powers of cubic fourfolds.* Trans. Amer. Math. Soc. **353** (2001), no. 4, 1455–1468